

Comment on “Geometric phase of entangled spin pairs in a magnetic field”

Yang Xiang* and Shi-Jie Xiong

*National Laboratory of Solid State Microstructures and Department of Physics,
Nanjing University, Nanjing 210093, China*

(Dated: February 1, 2008)

The degree of entanglement between two spins may change due to interaction. About this we find that a wrong result in a recent work by Ge and Wadati [Phys. Rev. A **72**, 052101(2005)] which breach the basic principle.

PACS numbers: 03.65.Ud, 03.65.Vf

Recently, Ge and Wadati [1] studied the geometric phase in the case of two initially entangled interacting $\frac{1}{2}$ -spins under a time-independent magnetic field. The Hamiltonian operator of the model is

$$H = \sum_{i=1,2} \omega_i S_z^{(i)} + \frac{8\lambda}{\hbar} \sum_{q=x,y,z} a_q S_q^{(1)} S_q^{(2)}, \quad (1)$$

where $\mathbf{S}^{(1)} = (S_x^{(1)}, S_y^{(1)}, S_z^{(1)})$ and $\mathbf{S}^{(2)} = (S_x^{(2)}, S_y^{(2)}, S_z^{(2)})$ are the spin operators of the two particles, ω_1 and ω_2 are the corresponding Larmor frequencies originating from a time-independent magnetic field along the z axis, λ and (a_x, a_y, a_z) are the strength and anisotropy of the interaction. There is an interaction between the two spins, so the degree of entanglement may change due to the interaction. But we find that Ge and Wadati [1] have ignored this.

Firstly, let us consider a $\frac{1}{2}$ -spin precessing in a time-independent uniform magnetic field pointing in the z direction. Assuming that the angle between the spin of the initial state $|\mathbf{n}(t=0)\rangle$ and the z axis is θ , the state at a later time t is

$$|\mathbf{n}(t)\rangle = e^{-i\varphi(t)/2} \cos\left(\frac{\theta}{2}\right) |+\rangle + e^{i\varphi(t)/2} \sin\left(\frac{\theta}{2}\right) |-\rangle. \quad (2)$$

Here $\varphi(t) = \varphi(0) + \omega t$ with $\varphi(0)$ being the initial azimuth angle with the x axis and ω is the Larmor frequency. At the same time, any pure state of two $\frac{1}{2}$ -spins can be decomposed as

$$|\Psi\rangle = e^{-i\beta/2} \cos\left(\frac{\alpha}{2}\right) |\mathbf{n}\rangle_1 |\mathbf{m}\rangle_2 + e^{i\beta/2} \sin\left(\frac{\alpha}{2}\right) |-\mathbf{n}\rangle_1 |-\mathbf{m}\rangle_2 \quad (3)$$

according to the Schmidt theorem [2], where $\mathbf{n}$ and $\mathbf{m}$ are two points on the Poincaré sphere, and the subscripts denote spins 1 and 2 respectively. The “angle” α in Eq. (3) determines the degree of entanglement in the state. For a bipartite pure state, the measure of entanglement is the von Neumann entropy of the reduced state in either of the two parties [3]. For the state $|\Psi\rangle$ in Eq. (3) we can calculate its entanglement $E(|\Psi\rangle)$ as

$$E(|\Psi\rangle) = -\cos^2\left(\frac{\alpha}{2}\right) \log_2 \cos^2\left(\frac{\alpha}{2}\right) - \sin^2\left(\frac{\alpha}{2}\right) \log_2 \sin^2\left(\frac{\alpha}{2}\right). \quad (4)$$

In [1], Ge and Wadati assumed that the initial state of two spin- $\frac{1}{2}$ particles is

$$|\Psi(0)\rangle = e^{-i\beta/2} \cos\left(\frac{\alpha}{2}\right) |\mathbf{n}(0)\rangle_1 |\mathbf{m}(0)\rangle_2 + e^{i\beta/2} \sin\left(\frac{\alpha}{2}\right) |-\mathbf{n}(0)\rangle_1 |-\mathbf{m}(0)\rangle_2, \quad (5)$$

where $|\mathbf{n}(0)\rangle$ and $|\mathbf{m}(0)\rangle$ are

$$|\mathbf{n}(0)\rangle = e^{-i\varphi_1/2} \cos\left(\frac{\theta_1}{2}\right) |+\rangle + e^{i\varphi_1/2} \sin\left(\frac{\theta_1}{2}\right) |-\rangle$$

and

$$|\mathbf{m}(0)\rangle = e^{-i\varphi_2/2} \cos\left(\frac{\theta_2}{2}\right) |+\rangle + e^{i\varphi_2/2} \sin\left(\frac{\theta_2}{2}\right) |-\rangle$$

(see Eq.(12) and Eq.(13) in [1]), then they argued that by applying $U(t) = e^{-iHt}$ on $|\Psi(0)\rangle$ they can obtain

$$|\Psi(t)\rangle = e^{-i\beta/2} \cos\left(\frac{\alpha}{2}\right) |\mathbf{n}(t)\rangle_1 |\mathbf{m}(t)\rangle_2 + e^{i\beta/2} \sin\left(\frac{\alpha}{2}\right) |-\mathbf{n}(t)\rangle_1 |-\mathbf{m}(t)\rangle_2 \quad (6)$$

(see Eq. (14) in [1]), where $|\mathbf{n}(t)\rangle$ and $|\mathbf{m}(t)\rangle$ are given by Eq. (2), and H is expressed in Eq. (1). It is noted that $|\pm\mathbf{n}(t)\rangle$ are always orthogonal with each other. As the “angle” α in the Schimdt decomposition determines the degree of entanglement in the state, from the comparison of Eq. (5) with Eq. (6), one can easily find that

$$E(|\Psi(0)\rangle) = E(|\Psi(t)\rangle), \quad (7)$$

i.e., the entanglement of $|\Psi(0)\rangle$ equals the entanglement of $|\Psi(t)\rangle$. However, this is not true when there exists an interaction between the two spins.

In fact, Sjöqvist [4] has studied the geometric phase of a closed quantal system consisting of two spin- $\frac{1}{2}$ particles with a spin-spin interaction. The Hamiltonian operator is

$$H = \frac{2\lambda}{\hbar} \mathbf{S}^{(1)} \cdot \mathbf{S}^{(2)}, \quad (8)$$

similar to the interaction term in Eq. (1). In this work Sjöqvist pointed out that the degree of entanglement may change due to the interaction between the two spins, a

product state may evolve into an entangled state and vice versa, and it is impossible to make one of the subsystems evolve but to keep the other unchanged. A simple example has been proposed to illustrate this in [4]. For two spin- $\frac{1}{2}$ particles under Hamiltonian of Eq. (8), if the initial state is

$$|\Psi(0)\rangle = \cos\left(\frac{a}{2}\right) |1, 0\rangle + \sin\left(\frac{a}{2}\right) |0, 0\rangle, \quad (9)$$

where $|1, 0\rangle = \frac{1}{\sqrt{2}}(|+\rangle_1 |-\rangle_2 + |-\rangle_1 |+\rangle_2)$ and $|0, 0\rangle = \frac{1}{\sqrt{2}}(|+\rangle_1 |-\rangle_2 - |-\rangle_1 |+\rangle_2)$, with subscripts denoting spin 1 and 2, we can express $|\Psi(0)\rangle$ in terms of the Schmidt decomposition

$$|\Psi(0)\rangle = \cos\left(\frac{\alpha}{2}\right) |+\rangle_1 |-\rangle_2 + \sin\left(\frac{\alpha}{2}\right) |-\rangle_1 |+\rangle_2, \quad (10)$$

where α satisfies $\tan(\frac{\alpha}{2}) = (\cos(\frac{a}{2}) - \sin(\frac{a}{2})) / (\cos(\frac{a}{2}) + \sin(\frac{a}{2}))$. This means that if the initial state is given, α is also given as a parameter. The value of α , however, may evolve in time under the interaction. By applying $U(t) = e^{-iHt}$ on $|\Psi(0)\rangle$, we can obtain $|\Psi(t)\rangle$ as

$$\begin{aligned} |\Psi(t)\rangle &= e^{-i\lambda t/2} \cos\frac{a}{2} |1, 0\rangle + e^{i3\lambda t/2} \sin\frac{a}{2} |0, 0\rangle \\ &= \frac{e^{i\lambda t/2}}{\sqrt{2}} \left[\left(e^{-i\lambda t} \cos\frac{a}{2} + e^{i\lambda t} \sin\frac{a}{2} \right) |+\rangle_1 |-\rangle_2 \right. \\ &\quad \left. + \left(e^{-i\lambda t} \cos\frac{a}{2} - e^{i\lambda t} \sin\frac{a}{2} \right) |-\rangle_1 |+\rangle_2 \right]. \end{aligned} \quad (11)$$

Ignoring an overall phase we have

$$|\Psi(t)\rangle = e^{-i\beta(t)/2} \cos\frac{\alpha(t)}{2} |+\rangle_1 |-\rangle_2$$

$$+ e^{i\beta(t)/2} \sin\frac{\alpha(t)}{2} |-\rangle_1 |+\rangle_2, \quad (12)$$

where $\cos\alpha(t) = \sin a \cos 2\lambda t$ and $\tan\beta(t) = -\tan a \sin 2\lambda t$. The “angle” $\alpha(t)$ is changing with time. This means that the degree of entanglement is also changing with time.

To summarize, we point out that there is a mistake in [1] which breaks a basic principle, i.e., the degree of entanglement may change due to the interaction between two spins. We illustrate this by using a simple example.

Acknowledgments This work was supported by the State Key Programs for Basic Research of China (2005CB623605 and 2006CB921803), and by National Foundation of Natural Science in China Grant Nos. 10474033 and 60676056.

* Electronic address: njuxy@sina.com

- [1] Xiang-Yu Ge and Miki Wadati, Phys. Rev. A **72**, 052101 (2005).
- [2] E. Schmidt, Ann. Math. **63**, 433 (1907).
- [3] C.H. Bennett, H.J Bernstein, S. Popescu, and B. Schumacher, Phys. Rev. A **53**, 2046 (1996).
- [4] Erik Sjöqvist, Phys. Rev. A **62**, 022109 (2000).